



Circulated as  
ISWI Newsletter Volume 2, Number 37

15 May 2010.



# Space Weather Research in Egypt

**A. Mahrous**



***Space Weather Monitoring Center, Faculty of Science, Helwan University, Cairo,  
Egypt. e-mail: [amahrous@helwan.edu.eg](mailto:amahrous@helwan.edu.eg), Fax.: 202-555-2468, Tel.: 202-556-7506***





# Outlines

- **Why we Study Space Weather ?**
- **First Space Weather Center in Egypt**
- **Geomagnetism Group**
- **Ionosphere Group**
- **Cosmic Ray Group**
- **Solar Physics Group**
- **Summary**

# Why we Study Space Weather ?



# Position Error




# Position Error




GPS Position Error

Ascension Island, 03/16/2002



# Spacecraft Damage/Loss



# Egypt is Located in Equatorial Anomaly Region (Crest and Trough)





Figure 1.5. Contour is the altitude profile of plasma density at 14LT, black lines are magnetic field lines and arrows stand for the directions of ion drifts [courtesy of Liu and Lin, 2006].

# Research Groups in our Center



# Our Journal publications

<b>Group</b>		<b>No of Res</b>	<b>Publications</b>
<b>Solar Physics</b>		<b>4</b>	<p><i>1) Empirical Model of the Transit Time of Interplanetary Coronal Mass Ejections</i> A. Mahrous, M. El-Nawawy, M. Hammama, and N. Ahmed, <i>Solar System Research</i>, 2009, Vol. 43, No. 2, pp. 128–135.</p> <p><i>2) CME–Fare Association During the 23rd Solar Cycle</i> A. Mahrous, M. Shaltout, M.M. Beheary, R. Mawad, M. Youssef <i>Advances in Space Research</i>, 2009, Vol. 43, pp. 1032–1035.</p>
<b>Ionosphere</b>		<b>5</b>	<p><i>Ionospheric Tomography Network of Egypt: A New Receiver Network in Support of the International Heliophysical Year</i> T. Garner, T. Gaussiran, J. York, D. Munton, C. Slack, A. Mahrous <i>Earth, Moon and Planet</i>, 2009, Vol. 104, pp. 227-235.</p>
<b>Cosmic Rays</b>		<b>3</b>	<p><i>Simulation of Muon-Induced Air Showers Affecting CMS Tracking Detectors</i> A. Mahrous, M. Sherif, and M. Soliman <i>Physics of Particles and Nuclei Letters</i>, 2009, Vol. 6, No. 3, pp. 246–250.</p>
<b>Geomagnetism</b>		<b>5</b>	<p><i>Mahrous, A., Ghamry, E., Elhawary, R., Fathy, I., Yamazaki, Y., Abe, S., Uozumi, T., Yumoto, K., First MAGDAS Installation at Fayum in Egypt</i>, <i>Advances in Space Research</i>, 2010, doi: 10.1016/j.asr.2010.04.022</p>

<http://www.helwan.edu.eg/english/Space>



# Space Weather Center



Home

About us

Members

Seminars

Projects

Research

## Space Weather Center



## Helwan University

A vertical sidebar containing several small profile pictures and associated text, likely representing the members or staff of the Space Weather Center.

# Joint Projects

Texas University (USA)  
CIDR Ionospheric Receiver

Kyushu University (Japan)  
MAGDAS Magnetometer

Stanford University (USA)  
AWESOME Ionospheric Receiver

SCINDA Ionospheric Receiver

European Union TEMPUS  
38,000 Euro (started)

US-Egyptian Joint Board  
180,000 US\$ (accepted)

Joining the African Network with  
European Networks (proposed)

Cyprus-Egyptian Joint Board  
90,000 EP (started)

# Geomagnetism Group

# MAGDAS Project



# MAGDAS Project


Prof. K. Yumoto, PI of MAGDAS project  
(during his visit to SWMC in 2008)



Associate Prof. Ayman Mahrous





## Installation of MAGDAS at FYM



# MAGDAS-II installation at ASW

Aswan, Egypt,  
15.20GMLat, 104.24G  
Installed at 08/12/23



# Typical Installation

- First, a solid foundation is laid for the sensor house. Then the sensor house is assembled with jumbo blocks. The sensor is accessible from the topside of the structure.





# Sensor Cable is Buried

The sensor cable is passed through a 5-cm flexible tube and then the tube is buried underground by about 10 cm.



Just prior to burial

# (MAGnetic Data Acquisition System/Circum-panPacific Magnetometer Network)



## MAGDAS-II Magnetometer

(W x D x H mm) or  
(φ x H mm)

CPMN-Amplifier



CPU BOX  
for Internet

(100x100x21 mm)



Internet  
(FTP or SSH)

Specified  
cable

(200x230x160 mm)

Analog

CPMN-Sensor



(200x200 mm)

Serial

A/D + Logger




(110x100x30 mm)

Data Acquisition  
Server



# Real-time Monitoring Data from FYM Station


ContentsViewer - playing




# Data is displayed as follows



# What we can get from MAGDAS




2000 : 7 : 15 : 00:00 -- 7 : 17 : 00:00 (Raw Data)





# Study of Sq (solar quiet) current globally



• We can see clear Sq current, equatorial electrojet, auroral electrojet, ring current and current connecting between northern and southern ionospheres on a disturbed day.

- This image was done by SERC staff.


# **Ionosphere Group**

**GPS Sub-group**

# GPS System at Helwan



# GPS System at Helwan



1: GPS receiver

2: GPS dual frequency antenna

3: Antenna cable (30 meter maximum)

4: Serial cable

5: Power cable

6: Personal computer running Linux


GPS dual frequency antenna




Helwan University







S4 AND ELEVATION ANGLE  
on November 24 2009




Scintillation Index(s4)




# TEC Profile



Vertical TEC (VTEC)



Vertical TEC (VTEC)



# Ionosphere Group


Simulation Sub-group

# Comparison with Simulations


**Thermospheric Ionospheric  
Electrodynamics General  
Circulation Model**

**TIEGCM**

# Simulation Results



**Energetic Event**



**Quite Day**

# Ionosphere Group

CIDR Sub-group


# CIDR Project 2008



# Coherent Ionospheric Doppler Receivers (CIDRs)




# Coherent Ionospheric Doppler Receivers (CIDRs)



- **7 Diamonds indicate CIDR systems**
- **Alaskan CIDRs (Stars) are owned by Univ. of Alaska-Fairbanks are part of a tomography chain with similar tomography receivers developed by NWRA (Triangles)**
- **Future deployments in New York and Columbia (Squares) have the equipment located at or near the site, but not running.**

# Coherent Ionospheric Doppler Receivers (CIDRs)



## Three CIDRs will be deployed to Egypt as part of IHY




- US coordinator (**Dr. Trevor Garner**), Texas University
- Egyptian coordinator (**Dr. Ayman Mahrous**), Helwan University.

The CIDR will be operated jointly by :

- 1- Helwan University
- 2- South Valley University
- 3- Alexandria University

# Coherent Ionospheric Doppler Receivers (CIDRs)

- Designed to track 150/400MHz LEO beacons (Transit/NIMS, GFO)
- Provides relative TEC and phase scintillation measurements at 50 Hz
- Useful for examining spatial structure with a relatively sparse receiver network and conducting ionospheric tomography



## **RADCAL/GFO Beacon Satellites**



**RADCAL (1993 to Present)**

- **Radio Altimetry and Ephemeris Satellites**
  - **150/400 MHz Radio Beacon**
  - **Ionospheric TEC Correction Data**



**GFO (1998 to Present)**



- **3 RADCAL/GFO Satellites**
- **20 RADCAL Ground Stations**
  - **Archived Data 1993 to Present**
  - **5 Second Samples**
  - **Maintained by AF Western Test Range Vandenberg**

**RADCAL on DMSP/F15  
(Aug 2006 to Present)**




# Advantages Over GPS

- **More accurate, no need for plasmaspheric corrections by using LEO satellites.**
- **Can measure the spatial structure of the ionosphere.**
- **A powerful tool for topographic image of the ionosphere**




# Event of October 11, 2008

A moderate magnetic storm was recorded on October 11, 2008 with a sudden commencement time occurred at 0838 UT. Figure shows, from top to bottom, the Kp index, Dst index = -52 nT, AE index, and IMF Bz component.



# Night-time: 2236 LT

Figure 8 shows the satellite track over Egypt (Oct 11, 2008 at 2036 UT, about 2236 LT at night time) is on the **recovery phase** ( $K_p=3$ ,  $Dst=-42$ ) and is an almost directly overhead pass by the satellite **OSCAR32**. The **TEC minimum** is located approximately at **30 degrees**. A mid-latitude trough also appeared at lower geographic latitudes, indicating that the trough is propagating equatorward as it is tracked in the three Fig.6,7 and 8 but all the result show pre-reversal enhancement at mid latitude. At latitudes closer to the magnetic equator, scintillations can also occur during nighttime. The scintillation is associated with spread-F occurrences. After local sunset, the bottom side of the F-region over the magnetic equator is subjected to gravitational Rayleigh-Taylor mechanisms. As a result, irregularities known as plasma bubbles are generated by rise to the topside ionosphere due to non-linear evolution of the instability and produce scintillations in discrete patches (Kumar and Gwal, 2000; Abdu et al., 1991).




# Cosmic Ray Group


Experimental Sub-group



# LHC



## The Large Hadron Collider in the LEP Tunnel



### Proton- Proton Collider

7 TeV + 7 TeV



Luminosity =  $10^{34} \text{cm}^{-2} \text{sec}^{-1}$

### first targets:





- Higgs boson (s)
- Supersymmetric Particles
- Quark-Gluon Plasma
- CP violation in B



A superconductive disk on the bottom, cooled by liquid nitrogen, causes the magnet above to levitate. The floating magnet induces a current, and therefore a magnetic field, in the superconductor, and the two magnetic fields repel to levitate the magnet.



# LHC Experiments





# What is the CMS experiment?





# CMS Outreach

37 Countries, 155 Institutes, 2000 scientists (including about 400 students)    October 2006

### TRIGGER, DATA ACQUISITION & OFFLINE COMPUTING

Austria, Brazil, CERN, Finland, France, Greece, Hungary, Ireland, Italy, Korea, Poland, Portugal, Switzerland, UK, USA

### TRACKER

Austria, Belgium, CERN, Finland, France, Germany, Italy, Japan\*, Mexico, New Zealand, Switzerland, UK, USA

### CRYSTAL ECAL

Belarus, CERN, China, Croatia, Cyprus, France, Italy, Japan\*, Portugal, Russia, Serbia, Switzerland, UK, USA

### PRESHOWER

Armenia, CERN, Greece, India, Russia, Taiwan

### RETURN YOKE

Barrel: Czech Rep., Estonia, Germany, Greece, Russia  
Endcap: Japan\*, USA

### SUPERCONDUCTING MAGNET

All countries in CMS contribute to Magnet financing in particular:  
Finland, France, Italy, Japan\*, Korea, Switzerland, USA

### FEET

Pakistan, China

### HCAL

Barrel: Bulgaria, India, Spain\*, USA  
Endcap: Belarus, Bulgaria, Georgia, Russia, Ukraine, Uzbekistan  
HO: India

### MUON CHAMBERS


Barrel: Austria, Bulgaria, CERN, China, Germany, Hungary, Italy, Spain  
Endcap: Belarus, Bulgaria, China, Colombia, Korea, Pakistan, Russia, USA

### FORWARD CALORIMETER

Hungary, Iran, Russia, Turkey, USA

Total weight : 12500 T  
Overall diameter : 15.0 m  
Overall length : 21.5 m  
Magnetic field : 4 Tesla

\* Only through industrial contracts



# Available Equipments



- 4 scintillators
- 4 scintillation detector boxes
- 4 Photo Multiplier Tubes PMT
- 4 electronic boxes to be attached to PMT
- Multichannel analyzer
- Digital oscilloscope
- High voltage power supply

# Cosmic Ray Group


Simulation Sub-group

# Muons Triggering to CMS

The interaction of cosmic ray particles in the upper atmosphere (primarily 9~15 Km above Earth's surface) usually produces pions (Duldig, 2000), a bound state of an up and anti-down quark.

With lifetime of ( $2.6 \times 10^{-8}$  s), the pion travels only hundreds of meters at velocities between (0.966 C and 0.977 C) before decaying into a muon and mu-neutrino .

The muons produced in that reaction descend to Earth's surface with ample supply of muons at sea level which facilitates the study of these particles (Caso et al., 2000).



# Data Analysis Group



**EUMED Grid**  
[empowering eScience across the Mediterranean]


HOME  
PUBLIC AREA

- ▶ partners
- ▶ applications
- ▶ work packages
- ▶ hosting a tutorial
- ▶ joining
- ▶ news
- ▶ press room
- ▶ press cuttings
- ▶ links
- ▶ contact us
- ▶ FAQs

- ▶ **country:** Egypt
- ▶ **author:** Prof. Mohamed Saleh
- ▶ **institute:** Helwan University
- ▶ **domain:** Bio-Informatics
- ▶ **contacts:**
- ▶ **description:** That application was a grid application running BLAST an algorithm for comparing primary biological sequence information (amino acid sequences of different proteins or the nucleotides of DNA sequence)
- ▶ **requirements:** The application requires BLAST software. It has been installed on EUMEDGRID e-Science Infrastructure and allowed CEs are

<http://www.eumedgrid.org/application/hero.html>

# EUMED Connect




# **Joint Projects with France, 2010**

## **(in progress)**

- **Micro satellite Programme for Solar and Space Weather Monitoring.**
- **GRID Project at the Space Weather Monitoring Center (SWMC).**
- **Monitoring of the water vapor in the troposphere along the River Nile.**

# Our Main Target : Space Weather (Monitoring & Prediction)



From Sun ..... to Earth

**Thank you**



**Space Weather Monitoring Center**