

題名 ISWI Newsletter – Vol. 3 No. 33
差出人 maeda@serc.kyushu-u.ac.jp

* ISWI Newsletter – Vol. 3 No. 33 28 March 2011 *
* *
* I S W I = International Space Weather Initiative *
* (www.iswi-secretariat.org) *
* *
* Publisher: Professor K. Yumoto, SERC, Kyushu University, Japan *
* Editor-in-Chief: Mr. George Maeda, SERC (maeda[at]serc.kyushu-u.ac.jp)*
* Archive location: www.iswi-secretariat.org (maintained by Bulgaria) *
* [click on "Publication" tab, then on "Newsletter Archive"] *
* Caveat: Under the Ground Rules of ISWI, if you use any material from *
* the ISWI Newsletter or Website, however minor it may seem *
* to you, you must give proper credit to the original source. *

Attachment(s):

(1) "Astrophysics noise -SW signal", 452 KB pdf, one page.

: Re:
: Astrophysics Noise:
: A Space Weather Signal
:

Dear ISWI Participant:

Attached is the first part of an article that may interest you.
EOS is the weekly publication of the AGU.

(The attached item is intentionally incomplete for copyright reasons.)

Faithfully yours,
: George Maeda
: The Editor
: ISWI Newsletter

EOS

EOS, TRANSACTIONS, AMERICAN GEOPHYSICAL UNION

This reprint is intentionally incomplete.



This pdf circulated in Volume 3, Number 33 on 28 March 2011.

IN THIS ISSUE: News: Earth Sciences Research Opportunities To Be Explored, p. 214
Meeting: Fifth Anniversary of the Landing on Titan, p. 215
About AGU: Historic Leadership Transition, p. 216
About AGU: Bibring Receives 2009 Whipple Award, p. 217
Research Spotlight: Exciting New Research From AGU Journals, p. 220

VOLUME 91 NUMBER 24 15 JUNE 2010

Astrophysics Noise: A Space Weather Signal

Imagine the accuracy of terrestrial weather forecasts if society relied on only a handful of isolated weather stations to supply all the input to meteorological models. Yet that is precisely the daunting situation faced by space weather forecasters, who seek to predict when and how ejections of plasma from the Sun will interact with the Earth's magnetosphere. These interactions can damage spacecraft electronics, produce spurious global positioning and navigation readings, interfere with radio communications, and disrupt electrical power line grids on the ground. Though modern society increasingly relies on satellite technology and electrical conveniences, only a handful of operating heliophysics missions supply the bulk of space weather model inputs.

Tremendous advances in the ability to anticipate and mitigate treacherous conditions at geosynchronous orbit and other

heavy ions such as carbon (C), oxygen (O), and iron (Fe) in charge states like C^{6+} , O^{7+} , and Fe^{13+} . Ionized close to the Sun, these species maintain their charge states as they flow outward with the solar wind through the solar system. When they encounter high densities of neutral atoms close to comets, the heavy ions undergo charge exchange, basically "stealing" an electron from the cometary neutral atoms. The resulting heavy ions, which now have a charge state lower by one, end up not in the ground state but rather in an excited state. This excited state quickly relaxes, causing the ions to emit soft X-ray photons. Since the original comet Hyakutake observations, scientists have learned that all comets are soft X-ray sources.

With the insight provided by the cometary soft X-ray observations, it is not much of a jump to realize that the same charge exchange process operating at comets must




Fig. 1. The correlation between soft X-ray intensities (red dashed curve) and solar wind particle fluxes (blue solid curve) shows that soft X-ray emissions near Earth respond strongly and quickly to changes in the solar wind.

