



Global Distribution of Quasi-Two-Day Waves: Results Based on the **NOGAPS-ALPHA Reanalysis Model**

D. Pancheva , P. Mukhtarov and D. Siskind

NIGGG, BAS, Sofia, Bulgaria

Naval Research Laboratory, Washington, USA



Motivations



- **QTDWs** are large-amplitude global-scale waves which can induce electric current systems in the ionosphere through a dynamo mechanism and in this way to be very efficient **direct dynamic agent in vertical coupling of the atmosphere-ionosphere system.**
- **QTDWs** effectively interact with **tides** generating new waves in the MLT region and having **additional impact on the ionosphere** through the secondary waves and modulated tides
- Although significant achievements have been attained in studying the spatial-temporal distribution of the QTDW phenomenon **there are still features hidden because of observational limitations.**



Used Data



- **Ground-based methods** - provide a detailed temporal variability of the QTDW however **cannot provide insight into its global distribution**
- **Satellite measurements** provides insight in the global spatial distribution however they **suffer from aliasing and poor temporal resolution problems.**
- **To overcome the above weak points we employ a version of the NOGAPS-ALPHA reanalysis model that provides output on an hourly basis.**

(Navy Operational Global Atmospheric Prediction System - Advanced Level Physics High Altitude version)

Data Analysis




- **14 months: 01 January 2009–28 February 2010**
- **All meteorological fields: neutral winds, temperature and geopotential height**
- **Altitudes between ~15 and ~95 km**
- **Latitudes between $\pm 80^\circ$**
- **2D Wavelet Analysis**
- **Decomposition procedure where QTDWS are separated simultaneously with tides and other PWs**




2D Wavelet Spectra

Eastward traveling QTDW

NH



SH







Mean 2D Wavelet Spectra

MLS/Aura GPH for 2005-2014




NH

SH




2D Wavelet Spectra over the Equator

GPH, U & T











Mean 2D Wavelet Spectra

MLS/Aura GPH for 2005-2014











E2 & E3 QTDW - Latitude Structure)



Day Number (start 01 January 2009)

E2 Altitude Structure











*hydrostatic
equation*


$$T = (H/R) \Phi_z$$

Sassi et al. (2002)


E3 Altitude Structure



E2 over the Equator - Altitude Structure





E2 over the Equator - Latitude Structure



E2 Altitude-Latitude Structure



$\lambda \sim 60 \text{ km}$




$\omega = Nk/m$
Kelvin wave



$\lambda \sim 30\text{-}35 \text{ km}$



E3 Altitude-Latitude Structure









2D Wavelet Spectra

Westward traveling QTDW

NH




SH



2D Wavelet Spectra over the Equator

Meridional Wind







Pancheva et al., (2006)



W2, W3 & W4 QTDW - Latitude Structure)

GPH

V




Day Number (start 01 January 2009)




Day Number (start 01 January 2009)

W2 Altitude Structure

NH




SH






Day Numbre (start 01 January 2009)

W3 Altitude Structure

NH




SH






W4 Altitude Structure






NH



SH



W1 over Equator (V)




Rossby-gravity wave

$\lambda \sim 25$ km


W2 over Equator & Middle Latitudes (V)

Rossby-gravity


V Amp 2d W2 PW (h=88 km)




Amp 2-d W2 (360-380)




Amp 2-d W2 (180-210)



Ph 2-d W2 (360-380)



Ph 2-d W2 (180-210)




$\lambda \sim 65-70$ km

W3 & W4 Latitude-Altitude Structure (V)


W3

W4


Amp 2-d W3 (385-405)




Amp 2-d W3 (190-215)




Amp 2-d W4 (1-20)




Amp 2-d W4 (210-240)




Ph 2-d W3 (385-405)




Ph 2-d W3 (190-215)



Ph 2-d W4 (1-20)



Ph 2-d W4 (210-240)




Latitude (degree)

Latitude (degree)

Latitude (degree)

Latitude (degree)

~2-d E2 & W3 waves in GPH




New Approach for Searching Traveling PWs

- A low-pass filter from the 4th degree with a boundary period of T_0 of 48 hours described as:

$$y(t) = \frac{1}{T} \int_{t-\frac{T_0}{2}}^{t+\frac{T_0}{2}} f(\tau) d\tau + \frac{2}{T} \sum_{k=1}^n \int_{t-\frac{T_0}{2}}^{t+\frac{T_0}{2}} f(\tau) \cos\left(\frac{2k\pi}{T}(t-\tau)\right) d\tau$$

with frequency characteristic:

$$H(T) = \sum_{k=-n}^n \frac{\sin\left(\pi\left(k - \frac{T_0}{T}\right)\right)}{\pi\left(k - \frac{T_0}{T}\right)}$$



Frequency Characteristics of Filter

Expansion in Fourier series as a function of longitude:

$$Y(lon, t) \approx Y_0(t) + \sum_{s=1}^S Y_s(t) \cos(slon - \varphi_s(t)) \quad (1)$$

Angular frequency is defined as the derivative of the phase:

$$\begin{aligned} \omega(t)t &= \varphi(t) \\ \frac{d\omega(t)}{dt}t + \omega(t) &= \frac{d\varphi(t)}{dt} \end{aligned} \quad \omega(t) = \frac{d\varphi(t)}{dt} \quad (2)$$


Having in mind (2) then (1) becomes:

$$Y(lon, t) \approx Y_0(t) + \sum_{s=1}^S Y_s(t) \cos(\omega_s(t)t - slon)$$




Examples

GPH 2-d E2 wave (SH)




GPH 2-d E3 wave (NH)





Example V over Equator

V 2-d W1 wave (eq)



Instantaneous Frequency



Summary (new 2-d waves)



- Two different types of eastward-traveling waves are identified: (i) E2 & E3 waves at middle and high latitudes, observed in the winters, and (ii) E2 wave observed over the equator with not well defined seasonal variability but with some enhancement between June and August; most probably it is a Kelvin wave.
- 2-d E2 wave is a primary wave, i.e. it is not generated by coupling between DW1 and 2-d W3 wave
- W1 & W2 waves Rossby-gravity waves observed in the meridional wind at low latitudes

THANK YOU

